



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$+ \frac{c^2(x^2 + y^2)(b^4x^2 + a^4y^2)}{a^2b^2(a^2b^2 - b^2x^2 - a^2y^2)}]^{\frac{1}{2}} dx dy.$$

This expression does not seem to be easy to integrate.

MISCELLANEOUS.

96. Proposed by H. M. CASH, Lore City, Guernsey County, Ohio.

A stick of timber is 12 feet long, 8 inches deep, and 3 inches wide at one end, and 5 inches deep, and 12 inches wide at the other end. An what distance from either end should it be cut to divide it into two equal parts?

Solution by H. C. WHITAKER, A. M., Ph. D., Manual Training School, Philadelphia, Pa., and G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa.

By the Prismoidal Formula, Volume = $\frac{1}{6}(\text{upper base} + \text{lower base} + 4 \text{ middle section}) \times \text{height}$.

Hence, the volume of the whole stick equals

$$\frac{1}{6}(8 \times 3 + 5 \times 12 + 4 \times \frac{1}{2} \times \frac{1}{2}) \times 144 = 6696.$$

Denote the distance of the required section from the 8×3 end by x , the depth at that section by y , the width by z . Then $y = 8 - \frac{1}{8}x$, and $z = 3 + \frac{1}{8}x$; also

$$\frac{1}{6} \left[8 \times 3 + yz + 4 \left(\frac{8+y}{2} \right) \left(\frac{3+z}{2} \right) \right] x = 3348.$$

$$\text{Whence } x^3 - 504x^2 - 55296x + 7713792 = 0.$$

$$x = 84.8856 \text{ inches} = 7 \text{ feet, } 0.8856 \text{ inches.}$$

Also solved by R. L. MOORE.

97. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

A spherical soap-bubble is electrified in such a way that the excess of the internal over the external air pressure is 2π when the bubble is in equilibrium. How does the tension of the film vary with the electric density?

Solution by the PROPOSER.

Let p be the excess of internal over external pressure, σ = electrical density, t = tension, r = radius of bubble.

Then $p + 2\pi\sigma^2 = 2t/r$, (see Minchin, Vol. II, page 487, third edition).

But $p = 2\pi$.

$$\therefore 2\pi r(1 + \sigma^2) = 2t, \text{ or } \frac{t}{1 + \sigma^2} = \pi r.$$